

# Modeling Gastric Luminal Rifabutin Concentrations: RHB-105 (Rifabutin 50 mg Q8H) Provides More Favorable Exposure for H. pylori Eradication than Generic Rifabutin 150 mg BID or 300 mg QD

### Colin W. Howden<sup>1</sup>, Salil N. Pendse<sup>2</sup>, Mansi S. Jamindar<sup>3</sup>, Kely L. Sheldon<sup>3</sup>

<sup>1</sup>University of Tennessee College of Medicine, Memphis, TN; <sup>2</sup>Allucent, Durham, NC; <sup>3</sup>RedHill Biopharma, Medical Affairs, Raleigh, NC.


### BACKGROUND

- Helicobacter pylori (H. pylori) is the strongest risk factor for peptic ulcer disease and non-cardia gastric cancer, and has been linked to dyspepsia, iron deficiency anemia, and vitamin B12 deficiency.<sup>1-4</sup>
- Topical gastric antibiotic exposure is important for successful eradication of *H. pylori*.<sup>5,6</sup>
- Variable success has been reported with rifabutin when used off-label (typically dosed at 150 mg QD, 150 mg BID, or 300 mg QD) as a component of treatment.<sup>7</sup>
- In two phase 3 trials in treatment-naïve subjects, low dose rifabutin (50 mg Q8H) triple therapy, RHB-105, (Talicia<sup>®</sup>), (50 mg rifabutin, 1000 mg amoxicillin, 40 mg omeprazole Q8H for 14 days) demonstrated eradication rates of 89.4% (NCT03198507; modified intention-to-treat) and 83.8% ITT (NCT01980095; 90.3% in confirmed adherent subjects).<sup>8,9</sup>
- Previous work has demonstrated that RHB-105 (50 mg rifabutin Q8H) achieved intragastric rifabutin concentrations  $\geq$ MIC<sub>90</sub> for ~93% of the day compared to generic substitution (150 mg rifabutin QD) which only achieved rifabutin concentrations  $\geq$  MIC<sub>60</sub> for ~35% of the day.<sup>10</sup>
- Our aim was to use physiologically based pharmacokinetic (PBPK) modeling to assess how generic rifabutin dosing (150 mg BID and 300 mg QD) affects intragastric rifabutin concentrations and any potential impact on *H. pylori* eradication.

### METHODS

- Plasma pharmacokinetic (PK) data for rifabutin following 50 mg Q8H or 150 mg QD triple therapy were obtained from literature sources and clinical studies.
- Key chemical and biological properties for the rifabutin were obtained from literature sources or calculated using quantitative structure-activity relationship models.
- The remaining parameters were estimated by fitting model predictions to a subset of the plasma PK data.
- The final parameterized PBPK model was validated against the plasma PK data not used for fitting. The validated PBPK model was then used to simulate 150 mg BID, 300 mg QD, or 50 mg Q8H at steady state and assess both plasma and intragastric rifabutin concentrations (Figure 1).
- Simulated concentrations were used to predict time above the rifabutin  $MIC_{00}$  for *H. pylori* in the gastric lumen.
- Since RHB-105 includes omeprazole, the impact of intragastric pH on rifabutin absorption was explored along with the impact of rifabutin autoinduction on time above  $MIC_{00}$ .

#### Figure 1: Simplified Schematic of Rifabutin Transit Highlighting the Two Sites of Pharmacokinetic Modeling: (1) Intragastric Rifabutin Concentrations and (2) Plasma Rifabutin Concentrations



Created with Biorender.com

Rifabutin given orally must exit the stomach into the small intestine where it is absorbed and enters the blood stream (box 1; intragastric concentration) and is then re-secreted into the gastric lumen (box 2; plasma concentration).

## For more detailed information, please contact the Medical Affairs Team at RedHill Biopharma at: medinfo@redhillus.com

### RESULTS


rifabutin 150 mg QD (blue).

- - Plasma rifabutin concentrations comparing 50 mg Q8H vs. 150 mg QD were modeled with equivalent doses of omeprazole in each regimen (Figure 2). • Subsequently, time when intragastric concentration was  $\geq$  MIC<sub>90</sub> were modeled (Figure 3); time  $\geq$  MIC<sub>90</sub> was 22.25±1.08
  - hrs (93% of the day) for 50 mg Q8H vs.  $8.29 \pm 1.67$  hrs (35% of the day) for 150 mg QD (Table 1).

### Figure 2: Plasma Rifabutin Concentrations Comparing 50 mg Q8H (RHB-105) and 150 mg QD Over a 24-hour Period



Figure 3: Intragastric Rifabutin Concentrations Comparing 50 mg Q8H (RHB-105) and 150 mg QD Over a 24hour Period




Steady state mean local intragastric rifabutin concentrations over 24 hours modeled on day 6 of a 14-day regimen comparing 50 mg Q8H (orange) and 150 mg QD (blue).

#### Table 1: Time Intragastric Rifabutin Concentration $\geq$ MIC<sub>on</sub> with 50 mg Q8H vs. 150 mg QD


Regimens Parameters		Intragastric Rifabutin Concentrations: Time above MIC <sub>90</sub>	
		Without Meals	With Meals
Rifabutin 50 mg Q8H	Mean (SD), hr	10.85 (4.69)	22.25 (1.08)
	% Day	45.21	92.71
Rifabutin 150 mg QD	Mean (SD), hr	3.98 (2.68)	8.29 (1.67)
	% Day	16.6	34.54
Ratio of Time Intragastric Rif	abutin Concentration above		
Ratio of Time Intragastric Rifabutin Concentration above MIC <sub>90</sub> : 50 mg Q8H / 150 mg QD		2.73	2.68

Note: SD, Standard Deviation; Represents Day 6



and 300 mg QD (green).


150 mg BID, and 300 mg QD




• Although both generic regimens contain twice the total daily dose of rifabutin than RHB-105, both failed to achieve the prolonged time intragastric rifabutin concentration was  $\geq$ MIC<sub>oo</sub> (22.25±1.08 hrs; 93% of the day) seen with RHB-105 (50 mg Q8H).

#### Figure 4: Intragastric Rifabutin Concentrations Comparing 50 mg Q8H (RHB-105), 150 mg BID, and 300 mg QD Over a 24-hour Period

Steady state mean plasma rifabutin concentrations over 24 hours modeled on day 6 of a 14-day regimen comparing rifabutin 50 mg Q8H (orange) and





#### Table 2: Time Intragastric Rifabutin Concentration $\geq$ MIC<sub>60</sub> with 50 mg Q8H vs. 150 mg BID or 300 mg QD

Regimens Parameters		Intragastric Rifabutin Concentrations: Time above MIC <sub>90</sub>	
		Without Meals	With Meals
Rifabutin 50 mg Q8H*	Mean (SD), hr	10.85 (4.69)	22.25 (1.08)
	% Day	42.21	92.71
Rifabutin 150 mg BID**	Mean (SD), hr	7.24 (4.44)	16.32 (2.25)
	% Day	30.16	68.0
Rifabutin 300 mg QD**	Mean (SD), hr	3.96 (2.84)	8.51 (1.86)
	% Day	16.50	35.46

Note: SD, Standard Deviation \*40 mg omeprazole, given Q8H \*\*20 mg omeprazole, given BIDz



Steady state mean intragastric rifabutin concentrations over 24 hours modeled on day 6 of a 14-day regimen comparing 50 mg Q8H (orange), 150 mg BID (blue),

#### Figure 5. Proportion of Day (%) Intragastric Rifabutin Concentration is $\geq$ MIC<sub>on</sub> Comparing 50 mg Q8H (RHB-105),

### CONCLUSIONS

- Key determinants of intragastric rifabutin concentrations include gastrointestinal transit time, time required for systemic absorption, and dosing frequency.
- Rifabutin 50 mg Q8H (RHB-105) maintains intragastric rifabutin concentrations above its MIC<sub>oo</sub> for 3.12-fold longer than 150 mg QD.
- Here, we show that two other generically substituted rifabutin regimens (150 mg BID and 300 mg QD) fail to achieve and maintain intragastric rifabutin concentrations seen with rifabutin 50 mg Q8H (RHB-105) (Table 2) despite comprising twice the total daily dose of rifabutin.
- $\Box$  50 mg rifabutin Q8H 22.25±1.08 hrs (93% of the day).
- $\Box$  150 mg rifabutin BID 16.32±2.25 hrs (68% of the day).
- $\Box$  300 mg rifabutin QD 8.51±1.86 hrs (35% of the day).
- With 150 mg QD, 150 mg BID or 300 mg QD dosing, the shorter duration of time with intragastric rifabutin concentration above its MIC<sub>90</sub> for H. pylori could explain the lower eradication rates than seen with **RHB-105.**
- Moreover, generically substitued 150 mg BID and 300 mg QD rifabutin regimens fail to achieve intragastric concentrations seen with RHB-105, while doubling the daily dose of rifabutin, potentially exposing patients to serious dose-related adverse events.

### REFERENCES

- Adamu MA, Weck MN, Rothenbacher D, Brenner H. Incidence and risk factors for the development of chronic atrophic gastritis: five year follow-up of a population-based cohort study. Int J Cancer. 2011;128(7):1652-1658. https://doi.org/10.1002/ijc.25476
- EI-Serag HB, Kao JY, Kanwal F, et al. Houston Consensus Conference on Testing for Helicobacter pylori Infection in the United States Clin Gastroenterol Hepatol. 2018;16(7):992-1002 e1006. https://doi.org/10.1016/j.cgh.2018.03.013 Eslick GD, Lim LL, Byles JE, Xia HH, Talley NJ. Association of Helicobacter pylori infection with gastric carcinoma: a meta-analysis.
- *Am J Gastroenterol*. 1999;94(9):2373-2379. https://doi.org/10.1111/j.1572-0241.1999.01360.x Kato S. Matsukura N. Tsukada K. et al. Helicobacter pylori infection-negative gastric cancer in Japanese hospital patients: incidence
- and pathological characteristics. *Cancer Sci*. 2007;98(6):790-794. https://doi.org/10.1111/j.1349-7006.2007.00478.x Kimura K, Ido K Fau - Saifuku K, Saifuku K Fau - Taniguchi Y, et al. A 1-h topical therapy for the treatment of Helicobacter pylori
- infection. Am J Gastroenterol. 1995;90(1):60-63. https://www.ncbi.nlm.nih.gov/pubmed/7801950 Satoh K. Treatment of Helicobacter pylori infection by topical administration of antimicrobial agents. *Scand J Gastroenterol Suppl.* 1996;214(56):discussion 57-60. https://www.ncbi.nlm.nih.gov/pubmed/8722409
- Gisbert JP. Rifabutin for the Treatment of Helicobacter Pylori Infection: A Review. *Pathogens*. 2020;10(1). https://doi.org/10.3390/ pathogens10010015
- . Graham DY. Canaan Y. Maher J. Wiener G. Hulten KG. Kalfus IN. Rifabutin-Based Triple Therapy (RHB-105) for Helicobacter pylori Eradication: A Double-Blind, Randomized, Controlled Trial. Ann Intern Med. 2020;172(12):795-802. https://doi.org/10.7326/M19-
- Kalfus IN, Graham DY, Riff DS, Panas RM. Rifabutin-Containing Triple Therapy (RHB-105) for Eradication of Helicobacter pylori: Randomized ERADICATE Hp Trial. *Antibiotics* (Basel). 2020;9(10). https://doi.org/10.3390/antibiotics9100685
- 10. Howden CW, Pendse SN, Bush MA, Almenoff JS, Sheldon KL. S1417 RHB-105 Q8H Rifabutin Dosing Provides Favorable Exposure for Helicobacter pylori Eradication. American Journal of Gastroenterology. 2021;116(1):S650-S650. 10.14309/01.ajg.0000779200. 28165.4c

### **QR CODE**

Please use the provided QR Code to the right, for the online version of this poster.

